

# Multi-step Intranasal (IN) Naloxone

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Address \_\_\_\_\_ Date Rx \_\_\_\_\_

**Rx: 2mg/2 mL Luer-Jet™ Luer-Lock needleless naloxone syringe plus mucosal atomizer device (MAD-300)**

**Quantity: 2 syringes + 2 MAD-300 devices**

**Sig (for suspected opioid overdose):**

**Spray 1 ml (1/2 of syringe) into each nostril.**

**Repeat after 2- 3 minutes if no or minimal response.**

**Refills: 2**

**Prescriber \_\_\_\_\_**

**Signature \_\_\_\_\_**

# Intramuscular (IM) Naloxone

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Address \_\_\_\_\_ Date Rx \_\_\_\_\_

**Rx: single-use 0.4 mg / 1 mL naloxone vial plus  
3 mL syringe w/ 23-25 gauge 1 inch IM needle**

**Quantity: 2 vials + 2 IM syringe/needle**

**Sig (for suspected opioid overdose):**

**Inject 1 mL in shoulder or thigh.**

**Repeat after 2-3 minutes if no or minimal response.**

**Refills: 2**

**Prescriber \_\_\_\_\_**

**Signature \_\_\_\_\_**

# Intranasal (IN) Naloxone Device

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Address \_\_\_\_\_ Date Rx \_\_\_\_\_

**Rx: Narcan®: 1 pack of two 4 mg/0.1 mL intranasal devices**

**Quantity: 1 pack**

**Sig (for suspected opioid overdose):**

**Spray 0.1 mL into one nostril.**

**Repeat with second device into other nostril after 2-3 minutes if no or minimal response.**

**Refills: 2**

**Prescriber \_\_\_\_\_**

**Signature \_\_\_\_\_**

# Intramuscular (IM) Naloxone Auto-injector

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

Address \_\_\_\_\_ Date Rx \_\_\_\_\_

**Rx: Evzio™: 1 pack of two 0.4 mg/0.4 mL prefilled  
auto-injector devices with one trainer device**

**Quantity: 1 pack**

**Sig (for suspected opioid overdose):**

**Inject into outer thigh as directed by English voice prompt system.  
Place black side firmly on outer thigh and depress and hold for 5  
seconds.**

**Repeat with second device in 2-3 minutes if no or minimal response.**

**Refills: 2**

**Prescriber \_\_\_\_\_**

**Signature \_\_\_\_\_**